

AD-A146 078

ELECTRON COLLISION PROCESSES IN MOLECULAR GASES(U)  
NATIONAL BUREAU OF STANDARDS BOULDER CO QUANTUM PHYSICS  
DIV A V PHELPS 30 SEP 84 ARO-17703.3-PH MIPR-ARO-9-81

1/1

UNCLASSIFIED

F/G 20/8


NL



END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ARO 17703.3-PH

(2)

AD-A146 078

ELECTRON COLLISION PROCESSES IN MOLECULAR GASES

Final Report

A. V. Phelps

September 30, 1984

U. S. Army Research Office

8-82  
MIPR 8-82/17703-P

Quantum Physics Division  
U.S. Bureau of Standards

Approved for Public Release;  
Distribution Unlimited

DTIC  
ELECTE  
OCT 1 1984  
S B D

DTIC FILE COPY

84 09 25 076

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARO 17703.3-PH	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) Electron Collision Processes in Molecular Gases		5. TYPE OF REPORT & PERIOD COVERED FINAL, January 1, 1981 to September 30, 1984
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) A. V. Phelps		8. CONTRACT OR GRANT NUMBER(s) <del>NRK</del> ARO 8-82
9. PERFORMING ORGANIZATION NAME AND ADDRESS Quantum Physics Division U.S. Bureau of Standards Boulder, CO 80303		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE September 30, 1984
		13. NUMBER OF PAGES 4
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  NA		
18. SUPPLEMENTARY NOTES  The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Electrons, dissociation, hydrogen, oxygen, energy distribution, ionization, anisotropic scattering, excited atom		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The results obtained under this project include (a) the experimental determination of coefficients for the dissociation of molecules in electrical discharges, (b) the development of improved techniques for the prediction of the role of electron induced processes such as dissociation and ionization in discharges, and (c) the development of models of high pressure discharges.		

# FINAL REPORT

ARO PROPOSAL NUMBER: 17703-P

CONTRACT OR GRANT NUMBERS: MIPR ARO 9-81; MIPR ARO 8-82; MIPR ARO 139-83

NAME OF INSTITUTION: U.S. National Bureau of Standards, Boulder, CO 80303

AUTHOR OF REPORT: A. V. Phelps

## LIST OF PUBLICATIONS OR TECHNICAL REPORTS:

"Transport data for the modeling of electrical breakdown and discharges," in Electrical Breakdown and Discharges, ed. by E.E. Kunhardt and L.H. Leusen (Plenum, New York, 1981), NATO ASI Series 81a, p. 109-132.

"Effect of electrons produced by ionization on electron energy distributions," Phys. Rev. A27, 2858-2867 (1983), with S. Yoshida and L.C. Pitchford.

"Anisotropic scattering of electrons by N<sub>2</sub> and its effect on electron transport," Phys. Rev. A (to be submitted September 1984), with L.C. Pitchford.

"The role of highly excited atoms in high pressure electrical discharges," Phys. Rev. (in preparation), with P.A. Vicharelli.

"Dissociation of H<sub>2</sub> by low energy electrons," J. Chem. Phys. (in preparation), with M.A. Islam.

## PARTICIPATING SCIENTIFIC PERSONNEL:

Dr. A. V. Phelps (principal investigator)  
Dr. M. A. Islam (postdoctorate)  
Dr. L. C. Pitchford (postdoctorate)  
Dr. B. Stumpf (postdoctorate)  
Dr. P. A. Vicharelli (postdoctorate)  
Mr. J. M. Urbanowicz (undergraduate)

No degrees were earned by participating scientific personnel.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
<i>PER CALL JC</i>	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<i>A-1</i>	

#### STATEMENT OF PROBLEM:

The objectives of this research were a) to obtain the experimental data and develop the techniques required to predict the role of electron impact induced dissociation of molecules in electrical discharges such as high-power diffuse discharge switches, and b) to develop improved models of transient discharges in high pressure gases.

#### RESULTS:

The results obtained under this project between January 1, 1981 and ~~June~~ <sup>SEPT.</sup> 30, 1984 can be conveniently divided into a) the experimental determination of coefficients for the dissociation of molecules in electrical discharges, b) the development of improved techniques for the prediction of the role of electron induced processes such as dissociation and ionization in discharges, and c) the development of models of high pressure discharges.

##### (a) Experimental measurement of molecular dissociation.

We have developed and applied an improved discharge technique for the measurement of the dissociation coefficients for molecules by moderate energy electrons. In order to measure the density of atoms in the same volume as that in which they are produced, we measure the relative atom density and the time constant for loss of atoms using the atomic resonance absorption method. These measurements are made absolute by comparing the absorption by atoms produced by electron impact with the absorption by atoms produced with a known efficiency by argon metastables. The metastable density is determined by measurements of absorption in the near infrared using lines of measured spectral width.

The experimental results obtained with H<sub>2</sub>-Ar mixtures have been analyzed to obtain coefficients for electron induced dissociation of the H<sub>2</sub> and for

metastable excitation and ionization of the Ar. At the higher  $E/n$  the ionization coefficients are in good agreement with predictions of our Boltzmann code. Our metastable excitation coefficients are much lower than expected on the basis of published values for pure Ar, but scale satisfactorily with recent unpublished data by Tachibana. The derived total  $H_2$  dissociation coefficients are consistent with the lower metastable production rates. These results will be used to analyze our data for electron induced dissociation in pure  $H_2$  discharges. Attempts to apply this technique to  $O_2$ -Ar discharges in order to obtain dissociation coefficients for  $O_2$  have been unsuccessful thus far because of the difficulties in operating the pulsed discharge in  $O_2$ .

(b) Prediction of electron transport and reaction coefficients.

In order to analyze the results of the measurements of the coefficients for dissociation of molecules such as  $H_2$  and  $O_2$  and to use these results to predict the behavior of electrons in gases such as air, one must have reliable techniques available for solution of the Boltzmann equation for electrons. Accordingly, we have been investigating the accuracy of techniques for solution of this equation. Firstly, we examined the effects of including the two electrons produced by ionization on calculated electron energy distributions and transport coefficients. We found that as the electric to gas density ratio  $E/n$  was increased it was necessary to use increasingly accurate descriptions of the distribution in energy of the secondary electrons. More recently, we investigated the effects of including anisotropic electron scattering on calculated electron energy distributions and transport coefficients. We find that the effects of an isotropic scattering are small for low  $E/n$  provided that the momentum transfer cross

section is used for elastic scattering. At  $E/n$  above  $500 \times 10^{-21} \text{ V m}^2$  anisotropic scattering effects become noticeable, but are less important than proper allowance for anisotropy in the electron energy distribution. We have also derived an effective cross section for the resonant rotational excitation of  $\text{N}_2$  which is able to reproduce the Russian data on rapid heating of  $\text{N}_2$  in electron beam excited  $\text{N}_2$ .

(c) Models of high pressure, transient discharges.

The portion of this research carried out under ARO support was concerned with the development of simplified techniques for the modeling of the role of highly excited atoms in high pressure discharges. In particular, we have formulated a simplified model of the rates of ionization and recombination of highly excited atoms which includes the effects of both electron and neutral particle collisions with high excited atoms. The results of this model are in a form appropriate for inclusion in a hydrodynamic model of the time development of constricted discharges. Beginning on January 1, 1983 this research was transferred to AFOSR support.



**END**

**FILMED**

**10-84**

**DTIC**